arXiv:1602.07855v1 [math.GT] 25 Feb 2016

RIBBON-CLASP SURFACE-LINKS AND NORMAL FORMS OF SINGULAR SURFACE-LINKS

SEIICHI KAMADA AND KENGO KAWAMURA

Abstract. We introduce the notion of a ribbon-clasp surface-link, which is a generalization of a ribbon surface-link. We generalize the notion of a normal form on embedded surface-links to the case of immersed surface-links and prove that any (immersed) surface-link can be described in a normal form. It is known that an embedded surface-link is a ribbon surface-link if and only if it can be described in a symmetric normal form. We prove that an (immersed) surface-link is a ribbon-clasp surface-link if and only if it can be described in a symmetric normal form. We also introduce the notion of a ribbon-clasp normal form, which is a simpler version of a symmetric normal form.

1. Introduction

In this paper an *immerse surface-knot* or simply a *surface-knot* means a closed connected and oriented surface generically immersed in $\mathbb{R}^4$. When it is embedded, we also call it an *embedded surface-knot*. A *surface-link* is a disjoint union of surface-knots. Two surface-links are said to be *equivalent* if they are ambient isotopic.

A surface-link is called *trivial* or *unknotted* if it is the boundary of a disjoint union of embedded handlebodies in $\mathbb{R}^4$. A surface-link is called *ribbon* if it is the boundary of immersed handlebodies in $\mathbb{R}^4$ whose multiple point set is a union of ribbon singularities. (The definition of a ribbon singularity is given in Section 2. For an immersion $f : M \to \mathbb{R}^4$ of a compact 3-manifold M, the *boundary* of the immersed 3-manifold $f(M)$ means the image $f(\partial M)$ of the boundary ∂M of M.)

Definition 1.1. A surface-link is *ribbon-clasp* if it is the boundary of immersed handlebodies in $\mathbb{R}^4$ whose multiple point set is a union of ribbon singularities and clasp singularities.

2000 *Mathematics Subject Classification.* 57M25, 57M27, 57Q45.

Key words and phrases. Surface-link; ribbon-clasp surface-link; ribbon singularity; clasp singularity; normal form.

The definition of a clasp singularity is given in Section 2. By definition, a trivial surface-link is ribbon, and a ribbon surface-link is ribbon-clasp.

Let F be a ribbon-clasp surface-link and let $f : M \to \mathbb{R}^4$ be an immersion of handlebodies M with $F = f(\partial M)$ such that the multiple point set is a union of ribbon singularities and c (≥ 0) clasp singularities. There are two double points of F in each clasp singularity, and one of them is a positive double point and the other is negative. Thus F has c positive double points and c negative double points, and the normal Euler number of F is zero.

Theorem 1.2. *For a surface-link F, the following conditions are equivalent.*

(1) *F is a ribbon-clasp surface-link.*
(2) *F is obtained from a ribbon surface-link by finger moves.*
(3) *F is obtained from a trivial 2-link by 1-handle surgeries and finger moves.*
(4) *F is obtained from an M-trivial 2-link by 1-handle surgeries.*

1-handle surgeries, finger moves and M-trivial 2-links are explained in Section 2 and Theorem 1.2 is proved in Section 3.

In Section 4 we define a normal form for a ribbon-clasp surface-link called a *ribbon-clasp normal form* in terms of the motion picture method. We prove the following theorem in Section 5.

Theorem 1.3. *A surface-link is ribbon-clasp if and only if it is equivalent to a surface-link in a ribbon-clasp normal form.*

In Section 4 we also define a normal form for a (singular) surface-link. It is a generalization of a normal form for an embedded surface-link defined in Kawauchi-Shibuya-Suzuki [5].

Theorem 1.4. *Any surface-link F is equivalent to a surface-link F' in a normal form. Moreover, for any non-negative integers c'_+, c'_-, c''_+ and c''_- such that $c'_+ + c''_+$ is the number of positive double points of F and $c'_- + c''_-$ is the number of negative double points of F, we may assume that F' has c'_+ positive double points and c'_- negative double points in $\mathbb{R}^3 \times [0, \infty)$ and c''_+ positive double points and c''_- negative double points in $\mathbb{R}^3 \times (-\infty, 0]$.*

When F is an embedded surface-link, Theorem 1.4 is the theorem on normal forms given in [5] (cf. [4]).

We say that a surface-link is in a *symmetric normal form* if it is in a normal form and it is symmetric with respect to the hyperplane $\mathbb{R}^3 \times \{0\}$.

Theorem 1.5. *A surface-link is ribbon-clasp if and only if it is equivalent to a surface-link in a symmetric normal form.*

When F is an embedded surface-link, the above theorem is the characterization of a ribbon surface-link due to Kawauchi-Shibuya-Suzuki [6].

2. Preliminaries

In this section, we give definitions of a ribbon singularity, a clasp singularity, a 1-handle surgery, a finger move and an M-trivial 2-link.

Let M be a compact 3-manifold with non-empty boundary and $f : M \to \mathbb{R}^4$ an immersion of M into $\mathbb{R}^4$. Let Δ be a connected component of the multiple point set $\{x \in f(M) \mid \#f^{-1}(x) \geq 2\} \subset \mathbb{R}^4$.

We say that Δ is a *ribbon singularity* if Δ is a 2-disk in $\mathbb{R}^4$ and the preimage of Δ is the disjoint union of embedded 2-disks Δ_1 and Δ_2 in M such that Δ_1 is properly embedded in M and Δ_2 is embedded in the interior of M. Figure 1 shows a local model of a ribbon singularity and a motion picture.



Figure 1. A local model of a ribbon singularity

We say that Δ is a *clasp singularity* if Δ is a 2-disk in $\mathbb{R}^4$ and the preimage of Δ is the disjoint union of embedded 2-disks Δ_1 and Δ_2 in M such that for each $i \in \{1, 2\}$, $\partial\Delta_i$ is the union of two arcs α_i and β_i, where α_i is a properly embedded arc in M and β_i is a simple arc in ∂M which connects endpoints of α_i. Figure 2 shows a local model of a clasp singularity and a motion picture.



FIGURE 2. A local model of a clasp singularity

Let F be a surface-link. A *chord* attaching to F means an unoriented simple arc γ in $\mathbb{R}^4$ such that $F \cap \gamma = \partial\gamma$, which misses the double points of F. Two chords are *equivalent* or *homotopic* with respect to F if they are homotopic through chords attaching to F.

A 1-*handle* attaching to F means an embedded 3-disk B in $\mathbb{R}^4$ such that $F \cap B$ is the union of a pair of mutually disjoint 2-disks in ∂B, $F \cap B$ misses the double points of F, and the orientation of $F \cap B$ induced from ∂B is opposite to the orientation induced from F. Put

$$\mathrm{h}(F; B) := \mathrm{Cl}(F \cup \partial B - F \cap B),$$

which we call the surface-link obtained from F by a 1-*handle surgery* along B. (Here Cl means the closure.) Two 1-handles attaching to F are said to be *equivalent* if they are ambient isotopic by an isotopy of $\mathbb{R}^4$ keeping F setwise fixed. It is known that 1-handles attaching to F are equivalent if and only if their cores are equivalent as chords attaching to F. (Boyle [1] and Hosokawa-Kawauchi [3]). For a chord γ attaching to F, we denote by $\mathrm{h}(F; \gamma)$ the surface-link obtained from F by a 1-handle surgery along a 1-handle whose core is γ.

A finger move is the inverse operation of the Whitney trick (cf. [2, 7]). Here we use an alternative definition using a Montesinos twin and 1-handle surgeries. A *Montesinos twin* is a surface-link which is the boundary of a pair of embedded oriented 3-disks B_1^3 and B_2^3 with a single ribbon singularity between B_1^3 and B_2^3. It has a positive double

point and a negative double point. The equivalence class, as a surface-link, of a Montesinos twin is unique. We denote by T a Montesinos twin.

Let F be a surface-link. Let U be a 4-disk in $\mathbb{R}^4$ disjoint from F. Put a Montesinos twin $T = S_1 \cup S_2$ in U and let B_1^3 and B_2^3 be embedded 3-disks in U with a clasp singularity with $S_i = \partial B_i^3$ $(i \in \{1, 2\})$. Take a point $p_1 \in S_1 - S_1 \cap S_2$, a point $p_2 \in S_2 - S_1 \cap S_2$ and two distinct points q_1, q_2 of F missing the double points of F. Let γ_1 and γ_2 be oriented chords attaching to $F \cup M$ such that for $i \in \{1, 2\}$, γ_i starts from p_i and terminates at q_i and $\gamma_i \cap (B_1^3 \cup B_2^3) = \{p_i\}$. Let F' be the surface-link obtained from $F \cup T$ by 1-handle surgeries along two 1-handles whose cores are γ_1 and γ_2. Let γ be a chord attaching to F which is a concatenation of γ_1^{-1}, a simple arc from p_1 to p_2 in U and γ_2. We say that F' is obtained from F by a *finger move* along γ, which we denote by $\mathrm{f}(F; \gamma)$. It is seen that if γ and γ' are equivalent chords attaching to F then $\mathrm{f}(F; \gamma)$ is equivalent to $\mathrm{f}(F; \gamma')$.

Definition 2.1. An *M-trivial 2-link* is a surface-link which is a split union of a trivial 2-link and some (or no) Montesinos twins.

3. Proof of Theorem 1.2

Proof of Theorem 1.2. Suppose (1). Let F be a ribbon-clasp surface-link. Let $f : M \to \mathbb{R}^4$ be an immersion of handlebodies M with $F = f(\partial M)$ whose multiple point set is a union of ribbon singularities and clasp singularities. Let $\Delta_1, \ldots, \Delta_c$ be the clasp singularities, and for each $i \in \{1, \ldots, c\}$, let Δ_{i1} and Δ_{i2} be the 2-disks in M with $f^{-1}(\Delta_i) = \Delta_{i1} \cup \Delta_{i2}$. Let N_{ij} $(i \in \{1, \ldots, c\}, j \in \{1, 2\})$ be a regular neighborhood of Δ_{ij} in M and let D_{ij} be the properly embedded 2-disk in M with $D_{ij} = \mathrm{Cl}(\partial N_{ij} \cap \mathrm{Int} M)$. Let $D_{ij} \times [0, 1]$ be a regular neighborhood of D_{ij} in $\mathrm{Cl}(M - N_{ij})$. Put $M_0 = \mathrm{Cl}(M - \cup_{i,j}(N_{ij} \cup D_{ij} \times [0, 1])$ where i and j run over all $i \in \{1, \ldots, c\}$ and $j \in \{1, 2\}$. Let $f_0 : M_0 \to \mathbb{R}^4$ be the restriction of f to M_0. Since M_0 is a 3-manifold homeomorphic to M, which is a union of handlebodies. The multiple point set of the immersion f_0 is a union of ribbon singularities, and $F_0 := f_0(\partial M_0)$ is a ribbon surface-link. For each $i \in \{1, \ldots, c\}$, let $T_i := f(\partial N_{i1} \cup \partial N_{i2})$, which is a Montesinos twin. The image $f(D_{ij} \times [0, 1])$ is a 1-handle attaching to $F_0 \cup (\cup_i T_i)$, and we see that F is obtained from F_0 by finger moves using $T_1, \ldots, T_c$ and 1-handles $f(D_{ij} \times [0, 1])$. Thus we have (2).

Suppose (2). Let F be obtained from a ribbon surface-link F_0 by finger moves. The ribbon surface-link F_0 is obtained from a trivial 2-link, say F_{00}, by 1-handle surgeries. More precisely, there are mutually

disjoint 1-handles, say $B_1, \ldots, B_n$, attaching to F_{00} and the surgery result is F_0. If necessary modifying the finger moves, we may assume that the chords attaching to F_0 along which the finger moves are applied are disjoint from the 1-handles $B_1, \ldots, B_n$. Then we can apply the 1-handle surgeries along $B_1, \ldots, B_n$ and the finger moves simultaneously to F_{00}. Thus we have (3). Since $F_{00} \cup (\cup_i T_i)$ is an M-trivial 2-link, we also have (4).

Suppose (3). Let F be obtained from a trivial 2-link by 1-handle surgeries and finger moves. Without loss of generality, we may assume that F is obtained from a trivial 2-link by 1-handle surgeries first and then by finger moves, namely, F is obtained from a ribbon surface-link F_0 by finger moves. Thus we have (2).

Suppose (4). Let F be obtained from an M-trivial 2-link F' by 1-handle surgeries. There is an immersion $f' : M' \to \mathbb{R}^4$ with $f'(\partial M') = F'$ such that M' is a disjoint union of 3-disks and the multiple point set of f' is a union of clasp singularities. Modifying the cores of the 1-handles attaching to F' used to obtain F, we assume that the cores intersect with $f'(M')$ in general position. Then we have an immersion $f : M \to \mathbb{R}^4$ with $f(\partial M) = F$ which is an extension of f' such that M is a disjoint union of handlebodies and the multiple point set of f is a union of clasp singularities and ribbon singularities. Thus we have (1). This completes the proof of Theorem 1.2. □

4. Normal Forms

First we prepare some terminologies. We allow a link to be empty, which we call the empty link and denote it by $\emptyset$.

Let L be a link and let $\mathcal{D} = \{D_1, \ldots, D_m\}$ be a set of mutually disjoint m (≥ 1) oriented 2-disks in $\mathbb{R}^3$ with $L \cap D_i = \emptyset$ for $i \in \{1, \ldots, m\}$. We say that the link $L \cup (\cup_i \partial D_i)$ is obtained from L by a 0-*surgery* along $\mathcal{D}$. We also say that L is obtained from $L \cup (\cup_i \partial D_i)$ by a 2-*surgery*.

Let L be a link and let $\mathcal{D} = \{D_1, \ldots, D_m\}$ be a set of mutually disjoint oriented 2-disks in $\mathbb{R}^3$ such that for $i \in \{1, \ldots, m\}$, D_i is a band attaching to L. We say that the link

$$\mathrm{h}(L; \mathcal{D}) := \mathrm{Cl}(L \cup (\cup_i \partial D_i) - L \cap (\cup_i \partial D_i))$$

is obtained from L by a 1-*surgery*. A 1-surgery is also called *surgery* or *band surgeries*. (A *band* attaching to L is an oriented 2-disk D in $\mathbb{R}^3$ such that $L \cap D$ is the union of a pair of mutually disjoint arcs in ∂D and the orientation of $L \cap \partial D$ induced from ∂D is opposite to the orientation induced from L.)

Let L be a link and let $(h_s \mid s \in [0, 1])$ be an ambient isotopy of $\mathbb{R}^3$, i.e., each $h_s : \mathbb{R}^3 \to \mathbb{R}^3$ is a homeomorphism and $h_0 = \mathrm{id}$.

When $L' = h_1(L)$, we say that L' is obtained from L by an *isotopic deformation*. When each h_s is the identity map, we call the isotopic deformation the *identity deformation*.

Let L be a link and let L' be obtained from L by applying some crossing changes. There is a homotopy $(g_s : M^1 \to \mathbb{R}^3 \mid s \in [0,1])$ of the source circles M^1 of the link into $\mathbb{R}^3$ with $g_0(M^1) = L$ and $g_1(M^1) = L'$ such that each g_s, except $s = 1/2$, is an embedding of M^1 and at $s = 1/2$ intersections occur. We call such a homotopy a *crossing change deformation*. It is called a *simple crossing change deformation* if exactly one crossing change occurs.

A *link transformation* from L to L', denoted by $L \to L'$, means one of the following transformation changing L to L':

(1) an isotopic deformation by an ambient isotopy $(h_s \mid s \in [0,1])$ of $\mathbb{R}^3$,
(2) a k-surgery along $\mathcal{D}$ for some $k \in \{0,1,2\}$,
(3) a crossing change deformation by a homotopy $(g_s \mid s \in [0,1])$.

For an interval $[a,b]$ $(a < b)$, the *realizing surface* of $L \to L'$ in $\mathbb{R}^3 \times [a,b]$ is a compact oriented surface, say F, such that

(1) for an isotopic deformation by an ambient isotopy $(h_s \mid s \in [0,1])$ of $\mathbb{R}^3$, $F \cap \mathbb{R}^3 \times \{t\} = h_s(L) \times \{t\}$ for $t \in [a,b]$, where $s = (t-a)/(b-a)$,
(2) for a k-surgery along $\mathcal{D} = \{D_1, \ldots, D_m\}$ for some $k \in \{0,1,2\}$,

$$F \cap \mathbb{R}^3 \times \{t\} = \begin{cases} \mathrm{h}(L;\mathcal{D}) \times \{t\} & \text{for } m \in ((a+b)/2, b] \\ L \cup (\cup_i D_i) \times \{t\} & \text{for } m = (a+b)/2 \\ L \times \{t\} & \text{for } m \in [a, (a+b)/2), \end{cases}$$

(3) for a crossing change deformation by a homotopy $(g_s \mid s \in [0,1])$, $F \cap \mathbb{R}^3 \times \{t\} = g_s(L) \times \{t\}$ for $t \in [a,b]$, where $s = (t-a)/(b-a)$.

We denote the realizing surface by $F(L \to L')_{[a,b]}$.

A *link transformation sequence* is a sequence of link transformations $\mathcal{L} : L_1 \to L_2 \to \cdots \to L_m$. A *division* of an interval $[a,b]$ of length m is a sequence of real numbers $t_1, t_2, \ldots, t_m$ with $a = t_1 < t_2 < \cdots < t_m = b$. The *realizing surface* of a link transformation sequence $\mathcal{L} : L_1 \to L_2 \to \cdots \to L_m$ in $\mathbb{R}^3 \times [a,b]$ with a division $t_1, t_2, \ldots, t_m$ is the union of the realizing surfaces $F(L_i \to L_{i+1})_{[t_i, t_{i+1}]}$ for $i = 1, \ldots, m-1$. It is a surface in $\mathbb{R}^3 \times [a,b]$, which we denote by $F(\mathcal{L})_{[t_1,\ldots,t_m]}$ or $F(\mathcal{L})_{[a,b]}$.

We call a link L an *H-trivial link* with m Hopf links if it is a split union of a trivial link and m Hopf links for some $m \geq 0$.

We say that a link transformation $L \to L'$ is a *Hopf-splitting deformation* when $L \to L'$ is a crossing change deformation such that each crossing change occurs for a Hopf link to change it into a trivial 2-component link. A *simple Hopf-splitting deformation* is a Hopf-splitting deformation that changes exactly one Hopf link to a trivial 2-component link.

Note that when $L \to L'$ is a *Hopf-splitting deformation* from an H-trivial link L with m Hopf links to a trivial link L', the number of crossing changes is m.

Let $L \to L'$ be a link transformation that is a 1-surgery along $\mathcal{D} = \{D_1, \dots, D_m\}$. When $\#L' = \#L - m$, we call it a 1-surgery of *fusion type*. When $\#L' = \#L + m$, we call it a 1-surgery of *fission type*.

Definition 4.1. A surface-link F is in a *ribbon-clasp normal form* if it is the realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O \to L \to L_0 \to L \to O \to \emptyset$$

in $\mathbb{R}^3 \times [-3, 3]$ with a division $-3, -2, -1, 0, 1, 2, 3$ satisfying the following conditions.

(1) O is a trivial link and $O \to \emptyset$ is a 2-surgery.
(2) L is an H-trivial link and $L \to O$ is a Hopf-splitting deformation.
(3) L_0 is a link and $L_0 \to L$ is a 1-surgery.
(4) The realizing surface is symmetric with respect to $\mathbb{R}^3 \times \{0\}$.

Example 4.2. Figure 3 shows a surface-knot in a ribbon-clasp normal form.



FIGURE 3. A surface-link in a ribbon-clasp normal form

The following is the definition of a *normal form* for a (singular) surface-link.

Definition 4.3. Let F be a surface-link with μ components and g total genus. We say that F is in a *normal form* if it is the realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O_- \to L_- \to K_- \to L_0 \to K_+ \to L_+ \to O_+ \to \emptyset$$

in $\mathbb{R}^3 \times [-4, 4]$ with a division $-4, -3, \cdots, 3, 4$ satisfying the following conditions.

(1) O_- and O_+ are trivial links, $\emptyset \to O_-$ is a 0-surgery and $O_+ \to \emptyset$ is a 2-surgery.
(2) L_- and L_+ are H-trivial links, $O_- \to L_-$ is the inverse of a Hopf-splitting deformation, and $L_+ \to O_+$ is a Hopf-splitting deformation.
(3) K_- and K_+ are links with μ components, $L_- \to K_-$ is a 1-surgery of fusion type and $K_+ \to L_+$ is a 1-surgery of fission type.
(4) L_0 is a link with $\mu + g$ components, $K_- \to L_0$ is a 1-surgery of fission type and $L_0 \to K_+$ is a 1-surgery of fusion type.
(5) Let $\mathcal{D}^{--}$ ($\mathcal{D}^-$, $\mathcal{D}^+$ or $\mathcal{D}^{++}$, resp.) be the set of bands used for the 1-surgery $L_- \to K_-$ ($K_- \to L_0$, $L_0 \to K_+$ or $K_+ \to L_+$, resp.). Then any band in $\mathcal{D}^{--}$ is disjoint from any band in $\mathcal{D}^-$, and any band in $\mathcal{D}^{++}$ is disjoint from any band in $\mathcal{D}^+$.

Let F be in a normal form. Let c'_+, c'_-, c''_+ and c''_- be non-negative integers such that F has c'_+ positive double points and c'_- negative double points in $\mathbb{R}^3 \times [0, \infty)$ and c''_+ positive double points and c''_- negative double points in $\mathbb{R}^3 \times (-\infty, 0]$. Then c'_+ is the number of negative Hopf links in L_+, c'_- is the number of positive Hopf links in L_+, c''_+ is the number of positive Hopf links in L_- and c''_- is the number of negative Hopf links in L_-.

Example 4.4. The surface-knot illustrated in Figure 4 is an immersed 2-sphere called a *negative standard singular 2-knot*, which we denote by S_-. It has a negative double point.



FIGURE 4. A negative standard singular 2-knot

(1) When we take $c'_- = 1$ and $c'_+ = c''_+ = c''_- = 0$, we have a surface-knot in a normal form illustrated in Figure 5 which is equivalent to S_-. It is a realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O_- \to L_- \to K_- \to L_0 \to K_+ \to L_+ \to O_+ \to \emptyset$$

such that $O_- = L_- = K_- = L_0 = K_+$ is a trivial knot, L_+ is a positive Hopf link, and O_+ is a trivial 2-component link.



FIGURE 5. A negative standard singular 2-knot in a normal form

(2) When we take $c''_- = 1$ and $c'_+ = c'_- = c''_+ = 0$, we have a surface-knot in a normal form illustrated in Figure 6 which is equivalent to S_-. It is a realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O_- \to L_- \to K_- \to L_0 \to K_+ \to L_+ \to O_+ \to \emptyset$$

such that O_- is a trivial 2-component link, L_- is a negative Hopf link, and $K_- = L_0 = K_+ = L_+ = O_+$ is a trivial knot.



FIGURE 6. A negative standard singular 2-knot in another normal form

Example 4.5. Figure 7 shows a Montesinos twin in a symmetric normal form, which is a realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O \to L \to K \to L_0 \to K \to L \to O \to \emptyset$$

such that O is a trivial 2-component link and $L = K = L_0$ is a positive Hopf link.



FIGURE 7. A Montesinos twin in a normal form

The surface-link in a symmetric normal form in Figure 7 has $c'_- = c''_+ = 1$ and $c'_+ = c''_- = 0$. Since a Montesinos twin is equivalent to its mirror image, by considering the mirror image in each cross section

in the motion picture of Figure 7, we have a Montesinos twin in a symmetric normal form with $c'_- = c''_+ = 0$ and $c'_+ = c''_- = 1$.

5. Proof of Theorems 1.3 and 1.5

Proof of Theorem 1.3. We show that a ribbon-clasp surface-link F is equivalent to a surface-link in a ribbon-clasp normal form. By Theorem 1.2, F is obtained from an M-trivial 2-link by 1-handle surgeries. The M-trivial 2-link is equivalent to a surface-link F' in a ribbon-clasp normal form which is the realizing surface of a link transformation sequence

$$\mathcal{L}' : \emptyset \to O' \to L' \to L' \to L' \to O' \to \emptyset$$

in $\mathbb{R}^3 \times [-3, 3]$ with division $-3, -2, -1, 0, 1, 2, 3$ such that $L' \to L' \to L'$ consists of the identity deformations. Thus, F is equivalent to a surface-link obtained from F' by 1-handle surgeries. Deforming the cores of the 1-handles up to equivalence, we may assume that the cores, say $\gamma_1, \ldots, \gamma_m$, are in $\mathbb{R}^3 \times \{0\}$. Let $D_1, \ldots, D_m$ be bands attaching to L' in $\mathbb{R}^3$ whose cores are $\tilde{\gamma}_1, \ldots, \tilde{\gamma}_m$, where $\tilde{\gamma}_i$ ($i \in \{1, \ldots, m\}$) is an arc in $\mathbb{R}^3$ with $\tilde{\gamma}_i \times \{0\} = \gamma_i$. Put $B_i := D_i \times [-1/2, 1/2]$, and we have 1-handles $B_1, \ldots, B_m$ attaching to F'. Since the 1-handles are equivalent to the original 1-handles attaching to F', the surface-link F is equivalent to the surface-link obtained from F' by 1-handle surgeries along $B_1, \ldots, B_m$, which is in a ribbon-clasp normal form.

We show that a surface-link F in a ribbon-clasp normal form is a ribbon-clasp surface-link. Suppose that F is the realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O \to L \to L_0 \to L \to O \to \emptyset$$

in $\mathbb{R}^3 \times [-3, 3]$ with division $-3, -2, -1, 0, 1, 2, 3$ satisfying the conditions of Definition 4.1. Let F' be the realizing surface of the link transformation sequence

$$\mathcal{L}' : \emptyset \to O \to L \to L \to L \to O \to \emptyset$$

in $\mathbb{R}^3 \times [-3, 3]$ with division $-3, -2, -1, 0, 1, 2, 3$ which is obtained from $\mathcal{L}$ by replacing $L \to L_0 \to L$ with the identity deformations $L \to L \to L$. Then F' is an M-trivial 2-link. Note that F is obtained from F' by 1-handles surgeries. (Let $\mathcal{D} = \{D_1, \ldots, D_m\}$ be the set of bands attaching to L along which the 1-surgery $L \to L_0$ is applied. Then $D_1 \times [-1/2, 1/2], \ldots, D_m \times [-1/2, 1/2]$ are 1-handles attaching to F', and F is obtained from F' by surgery along these 1-handles.) By Theorem 1.2, F is a ribbon-clasp surface-link. □

Proof of Theorem 1.5. First we show that a ribbon-clasp surface-link is equivalent to a surface-link in a symmetric normal form. Let F be a ribbon-clasp surface-link. By Theorem 1.3, we may assume that F is in a ribbon-clasp normal form such that it is the realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O \to L \to L \to L_0 \to L \to L \to O \to \emptyset$$

in $\mathbb{R}^3 \times [-4, 4]$ with division $-4, -3, -2, -1, 0, 1, 2, 3, 4$ satisfying the conditions of Definition 4.1 with the identity deformation $L \to L$. Let $\mathcal{D}$ be the set of bands attaching to $L \to L_0$ along which the 1-surgery $L \to L_0$ is applied. Let $F = F_1 \cup \cdots \cup F_\mu$, where F_i is the ith component of F. The link L is divided into μ sub-links $L_1, \ldots, L_\mu$ such that L_i is the cross-section of F_i at $t = -2$. Choose a subset $\mathcal{D}''$ of $\mathcal{D}$ such that when we put $K := \mathrm{h}(L; \mathcal{D}'')$, the 1-surgery $L \to K$ along $\mathcal{D}''$ is a 1-surgery of fusion type and the number of components of K is μ. Let $\mathcal{D}' = \mathcal{D} - \mathcal{D}''$. Let

$$\mathcal{L}' : \emptyset \to O \to L \to K \to L_0 \to K \to L \to O \to \emptyset$$

be the link transformation sequence obtained from $\mathcal{L}$ by replacing $L \to L \to L_0 \to L \to L$ by $L \to K \to L_0 \to K \to L$ where $L \to K$ is the 1-surgery along $\mathcal{D}''$ and $K \to L_0$ is the 1-surgery along $\mathcal{D}'$. By an argument in [5], we may assume that $K \to L_0$ is a 1-surgery of fission type and the number of components of L_0 is $\mu + g$. Then F is equivalent to the realizing surface of $\mathcal{D}''$ in $\mathbb{R}^3 \times [-4, 4]$ with division $-4, -3, -2, -1, 0, 1, 2, 3, 4$, which is in a symmetric normal form.

We show that a surface-link in a symmetric normal form is a ribbon-clasp surface-link. Let F be a surface-link, with μ components and g total genus, in a symmetric normal form, i.e., it is the realizing surface of a link transformation sequence

$$\mathcal{L} : \emptyset \to O \to L \to K \to L_0 \to K \to L \to O \to \emptyset$$

in $\mathbb{R}^3 \times [-4, 4]$ with division $-4, -3, -2, -1, 0, 1, 2, 3, 4$ satisfying the following conditions.

(1) O is a trivial link and $O \to \emptyset$ is a 2-surgery.
(2) L is an H-trivial link and $L \to O$ is a Hopf-splitting deformation.
(3) K is a link with μ components and $K \to L$ is a 1-surgery of fission type.
(4) L_0 is a link with $\mu + g$ components and $L_0 \to K$ is a 1-surgery of fusion type.

(5) Let $\mathcal{D}'$ (or $\mathcal{D}''$) be the set of bands along which the 1-surgery $L_0 \to K$ (or $K \to L$) is applied. Then any band of $\mathcal{D}''$ is disjoint from any band of $\mathcal{D}'$.
(6) The realizing surface is symmetric with respect to $\mathbb{R}^3 \times \{0\}$.

Put $\mathcal{D} := \mathcal{D}' \cup \mathcal{D}''$, which is a set of bands attaching to L_0 such that the surgery result is L. Let

$$\mathcal{L}' : \emptyset \to O \to L \to L \to L_0 \to L \to L \to O \to \emptyset$$

be the sequence obtained from $\mathcal{L}$ by replacing $L \to K \to L_0 \to K \to L$ with $L \to L \to L_0 \to L \to L$ where $L \to L$ is the identity deformation, $L_0 \to L$ is the 1-surgery along $\mathcal{D}$ and $L \to L_0$ is its reverse. Then F is equivalent to the realizing surface of $\mathcal{L}'$ in $\mathbb{R}^3 \times [-4, 4]$ with division $-4, -3, -2, -1, 0, 1, 2, 3, 4$. Finally, the surface-link is equivalent to a surface-link in a ribbon-clasp normal form. □

6. Proof of Theorem 1.4

First we prepare lemmas.

Let $L \to L'$ be a simple crossing change deformation and let $F(L \to L')_{[a,b]}$ be the realizing surface in $\mathbb{R}^3 \times [a, b]$. Let W be a small 3-disk in $\mathbb{R}^3$ such that $W \times [a, b] \cap F(L \to L')_{[a,b]} = \emptyset$.

Lemma 6.1. *In the above situation, there is a link transformation sequence $\mathcal{L} : L = L_0 \to L_1 \to L_2 \to L_3 \to L_4 = L'$ satisfying the following.*

(1) *$F(L \to L')_{[a,b]}$ is ambient isotopic to $F(\mathcal{L})_{[a,b]}$ by an isotopy of $\mathbb{R}^3 \times [a, b]$ rel $\mathbb{R}^3 \times \{a, b\}$.*
(2) *$L \to L_1$ is a 1-surgery along two bands, and L_1 is the split union of a link $\tilde{L}'$ and a Hopf link, where $\tilde{L}'$ is a link equivalent to L'.*
(3) *$L_1 \to L_2$ is an isotopic deformation moving $\tilde{L}'$ onto L' and moving the Hopf link into W so that L_2 is the split union of L' and a Hopf link in W,*
(4) *$L_2 \to L_3$ is a simple Hopf-splitting deformation that changes the Hopf link in W to a trivial 2-component link in W,*
(5) *$L_3 \to L_4$ is a 2-surgery that eliminates the trivial 2-component link in W.*

Proof. Let U be a 3-disk in $\mathbb{R}^3$ which is a regular neighborhood of the double point of the singular link appearing as the cross-section of $F(L \to L')_{[a,b]}$ at $t = (a + b)/2$. The motion picture in the top of Figure 8 shows the restriction of $F(L \to L')_{[a,b]}$ to $U \times [a, b]$ in a case of a negative double point. Replace it with the surface as shown in the second and the third lows of Figure 8, namely, which is the realizing

surface of a sequence $\mathcal{L}^{(0)} : L = L_0^{(0)} \to L_1^{(0)} \to L_2^{(0)} \to L_3^{(0)} \to L_4^{(0)} = L'$ satisfying the following.

(1) $F(L \to L')_{[a,b]}$ is ambient isotopic to $F(\mathcal{L}^{(0)})_{[a,b]}$ by an isotopy of $\mathbb{R}^3 \times [a,b]$ rel $\mathbb{R}^3 \times \{a,b\}$.
(2) $L \to L_1^{(0)}$ is a 1-surgery along two bands, and $L_1^{(0)}$ is the split union of a link $\tilde{L}'$ and a Hopf link $H^{(0)}$, where $\tilde{L}'$ is equivalent to L'.
(3) $L_1^{(0)} \to L_2^{(0)}$ is a simple crossing change deformation between the components of $H^{(0)}$. Let $O_2^{(0)}$ be the trivial 2-component link obtained from $H^{(0)}$. So $L_2^{(0)}$ is a split union of $\tilde{L}'$ and $O_2^{(0)}$.
(4) $L_2^{(0)} \to L_3^{(0)} = \tilde{L}'$ is a 2-surgery that eliminates $O_2^{(0)}$.
(5) $L_3^{(0)} \to L_4^{(0)}$ is an isotopy deformation.

We change the sequence $\mathcal{L}^{(0)}$ by inserting an isotopic deformation just after $L \to L_1^{(0)}$ such that a small 3-disk in $\mathbb{R}^3$ containing the Hopf link $H^{(0)}$ is moved into W. Then we obtain a desired sequence $\mathcal{L}$. □



FIGURE 8. Moving a double point

Let $\mathcal{L} : L_0 \to \cdots \to L_m$ be a link transformation sequence and let $F(\mathcal{L})_{[a,b]}$ be the realizing surface in $\mathbb{R}^3 \times [a,b]$. Suppose that for some n, $L_n \to L_{n+1}$ in $\mathcal{L}$ is a simple crossing change deformation. Let W be a small 3-disk in $\mathbb{R}^3$ such that $W \times [a,b] \cap F(\mathcal{L})_{[a,b]} = \emptyset$.

Lemma 6.2. *In the above situation, there is a link transformation sequence $\mathcal{L}^{(1)} : L_0^{(1)} \to \cdots \to L_{m+3}^{(1)}$ satisfying the following.*

(1) *$F(\mathcal{L})_{[a,b]}$ is ambient isotopic to $F(\mathcal{L}^{(1)})_{[a,b]}$ by an isotopy of $\mathbb{R}^3 \times [a,b]$ rel $\mathbb{R}^3 \times \{a,b\}$.*
(2) *For each $k \in \{1,\dots,n\}$, $L^{(1)}_{k-1} \to L^{(1)}_k$ is the same with $L_{k-1} \to L_k$.*
(3) *$L^{(1)}_n \to L^{(1)}_{n+1}$ is a 1-surgery along two bands, and $L^{(1)}_{n+1}$ is the split union of a link $\tilde{L}_{n+1}$ and a Hopf link, where $\tilde{L}_{n+1}$ is a link equivalent to L_{n+1}.*
(4) *$L^{(1)}_{n+1} \to L^{(1)}_{n+2}$ is an isotopic deformation moving $\tilde{L}_{n+1}$ onto L_{n+1} and moving the Hopf link into W so that $L^{(1)}_{n+2}$ is the split union of L_{n+1} and a Hopf link H in W.*
(5) *For each $k \in \{n+3,\dots,m+1\}$, $L^{(1)}_k$ is the split union of L_{k-1} and H, and the link transformation $L^{(1)}_{k-1} \to L^{(1)}_k$ is the same with $L_{k-2} \to L_{k-1}$ on L_{k-2} and the identity on H.*
(6) *$L^{(1)}_{m+1} \to L^{(1)}_{m+2}$ is a simple Hopf-splitting deformation that changes the Hopf link H to a trivial 2-component link in W.*
(7) *$L^{(1)}_{m+2} \to L^{(1)}_{m+3}$ is a 2-surgery that eliminates the trivial 2-component link in W. Then $L^{(1)}_{m+3} = L_m$.*

Proof. Applying Lemma 6.1 to $L_n \to L_{n+1}$, we see that there is a link transformation sequence $\mathcal{L}' : L'_0 \to \cdots \to L'_{m+3}$ satisfying the following.

(1) $F(\mathcal{L})_{[a,b]}$ is ambient isotopic to $F(\mathcal{L}')_{[a,b]}$ by an isotopy of $\mathbb{R}^3 \times [a,b]$ rel $\mathbb{R}^3 \times \{a,b\}$.
(2) For each $k \in \{1,\dots,n\}$, $L'_{k-1} \to L'_k$ is the same with $L_{k-1} \to L_k$.
(3) $L'_n \to L'_{n+1}$ is a 1-surgery along two bands, and L'_{n+1} is the split union of a link $\tilde{L}_{n+1}$ and a Hopf link, where $\tilde{L}_{n+1}$ is a link equivalent to L_{n+1}.
(4) $L'_{n+1} \to L'_{n+2}$ is an isotopic deformation moving $\tilde{L}_{n+1}$ onto L_{n+1} and moving the Hopf link into W so that L'_{n+2} is the split union of L_{n+1} and a Hopf link H in W.
(5) $L'_{n+2} \to L'_{n+3}$ is a simple Hopf-splitting deformation that changes the Hopf link H in W to a trivial 2-component link.
(6) $L'_{n+3} \to L'_{n+4}$ is a 2-surgery that eliminates the trivial 2-component link in W. Then $L'_{n+4} = L_{n+1}$.
(7) For $k \in \{n+5,\dots,m+3\}$, $L'_{k-1} \to L'_k$ is the same with $L_{k-4} \to L_{k-3}$.

By an ambient isotopy, we deform the realizing surface $F(\mathcal{L}')_{[a,b]}$ so that the simple Hopf-splitting deformation and the 2-surgery in the above (5) and (6) occur after the link transformations in (7). Then we have a desired link transformation sequence $\mathcal{L}^{(1)}$. □

Proof of Theorem 1.4. Let F be a surface-link. Divide the set of positive (or negative) double points of F into two groups, say G'_+ and G''_+ (or G'_- and G''_-), with $\#G'_+ = c'_+$ and $\#G''_+ = c''_+$ (or $\#G'_- = c'_-$ and $\#G''_- = c''_-$).

By a similar argument with [5], there is a link transformation sequence $\mathcal{L}^{(0)}$ such that a realizing surface of $\mathcal{L}^{(0)}$ is equivalent to F, each crossing change deformation is simple, there is no 0-surgery in $\mathcal{L}^{(0)}$ except the first link transformation, and there is no 2-surgery in $\mathcal{L}^{(0)}$ except the last link transformation. (Note that the first link transformation of $\mathcal{L}^{(0)}$ is a 0-surgery from the empty link to a trivial link, and the last link transformation is a 2-surgery from a trivial link to the empty link.)

Take a double point p_1 from $G'_+ \cup G'_-$ and let $L^{(0)}_{n_0} \to L^{(0)}_{n_0+1}$ be the simple crossing change deformation in $\mathcal{L}^{(0)}$ whose realizing surface contains p_1. Apply Lemma 6.2 to $\mathcal{L}^{(0)}$ with $L^{(0)}_{n_0} \to L^{(0)}_{n_0+1}$ and we have a link transformation sequence such that a realizing surface of the sequence is equivalent to F, each crossing change deformation is simple, and the last three link transformations are a simple Hopf-splitting deformation from an H-trivial link to a trivial link, a 2-surgery that eliminates the trivial 2-component link obtained by the simple Hopf-splitting deformation, and a 2-surgery from a trivial link to the empty link. Combine the last two link transformations into a link transformation which is a 2-surgery from a trivial link to the empty link. Let $\mathcal{L}^{(1)}$ be the link transformation sequence. Then $\mathcal{L}^{(1)}$ is a link transformation sequence such that a realizing surface is equivalent to F, each crossing change deformation is simple, and the last two link transformations are a simple Hopf-splitting deformation from an H-trivial link to a trivial link and a 2-surgery from the trivial link to the empty link.

Take a double point p_2 from $(G'_+ \cup G'_-) - \{p_1\}$ and let $L^{(1)}_{n_1} \to L^{(1)}_{n_1+1}$ be the simple crossing change deformation in $\mathcal{L}^{(1)}$ whose realizing surface contains p_2. By the same argument as above, we have a link transformation sequence $\mathcal{L}^{(2)}$ such that a realizing surface of $\mathcal{L}^{(2)}$ is equivalent to F, each crossing change deformation is simple, and the last three link transformations are a simple Hopf-splitting deformation from an H-trivial link to an H-trivial link, a simple Hopf-splitting deformation from the H-trivial link to a trivial link and a 2-surgery from the trivial link to the empty link.

Continue this procedure until we cover all double points belonging to $G'_+ \cup G'_-$, and we have a link transformation sequence such that a realizing surface of the sequence is equivalent to F, each crossing change deformation is simple, and the last $c'_+ + c'_- + 1$ link transformations are

$c'_+ + c'_-$ simple Hopf-splitting deformations from H-trivial links, and a 2-surgery from a trivial link to the empty link.

For this link transformation sequence, we apply a similar argument on double points belonging to $G''_+ \cup G''_-$. However, this time we move the double points forward in the time direction. Then we have a link transformation sequence such that a realizing surface of the sequence is equivalent to F, each crossing change deformation is simple, there is no 0-surgery in the sequence except the first link transformation, there is no 2-surgery in the sequence except the last link transformation, the first $c''_+ + c''_- + 1$ link transformations are a 0-surgery from the empty link to a trivial link and $c''_+ + c''_-$ inverses of simple Hopf-splitting deformations, and the last $c'_+ + c'_- + 1$ link transformations are $c'_+ + c'_-$ simple Hopf-splitting deformations and a 2-surgery from a trivial link to the empty link.

Combine the first $c''_+ + c''_-$ inverses of simple Hopf-splitting deformations to a Hopf-splitting deformation, and combine the last $c'_+ + c'_-$ simple Hopf-splitting deformations to a Hopf-splitting deformation. Then we have a link transformation sequence, say $\mathcal{L}' : \emptyset \to O'_- \to L'_- \to \cdots \to L'_+ \to O'_+ \to \emptyset$, where $L'_- \to \cdots \to L'_+$ is a sequence consisting of 1-surgeries, such that a realizing surface of the sequence $\mathcal{L}'$ is equivalent to F. Applying the argument in [5] to this sequence with $L'_- \to \cdots \to L'_+$, we obtain a sequence $\mathcal{L} : \emptyset \to O_- \to L_- \to K_- \to L_0 \to K_+ \to L_+ \to O_+ \to \emptyset$ as desired. □

References

[1] J. Boyle, *Classifying 1-handles attached to knotted surfaces*, Trans. Amer. Math. Soc. **306** (1988), 475–487.

[2] A. J. Casson, *Three lectures on new-infinite constructions in 4-dimensional manifolds*, In: À la Recherche de la Topologie Perdue, (Eds. L. Guillou and A. Marin), Progr. Math. **62**, pp. 201–214, Birkhäuser Boston, Boston, MA, 1986.

[3] F. Hosokawa and A. Kawauchi, *Proposals for unknotted surfaces in four-spaces*, Osaka J. Math. **16** (1979), 233–248.

[4] S. Kamada, *Braid and Knot Theory in Dimension Four*, Mathematical Surveys and Monographs, Vol. 95, American Mathematical Society, Providence, RI, 2002.

[5] A. Kawauchi, T. Shibuya and S. Suzuki, *Descriptions on surfaces in four-space, I: Normal forms*, Math. Sem. Notes, Kobe Univ. **10** (1982), 75–125.

[6] A. Kawauchi, T.Shibuya and S.Suzuki, *Descriptions on surfaces in four-space, II: Singularities and cross-sectional links*, Math. Sem. Notes, Kobe Univ. **11** (1983), 31–69.

[7] R. Kirby, *The topology of 4-manifolds*, Lecture Notes in Mathematics **1374**, Springer-Verlag, Berlin, 1989.

(S. Kamada) Department of Mathematics, Osaka City University, Sugimoto 3-3-138, Sumiyoshi-ku, Osaka 558-8585, Japan
E-mail address: skamada@sci.osaka-cu.ac.jp

(K. Kawamura) Department of Mathematics, Osaka City University, Sugimoto 3-3-138, Sumiyoshi-ku, Osaka 558-8585, Japan
E-mail address: k.kawamura0403@gmail.com